

## PE Human BCMA (CD269) Protein (C-Fc)

|                           |                            |
|---------------------------|----------------------------|
| <b>Catalog Number:</b>    | 802401, 802402             |
| <b>Size:</b>              | 25 ug, 100 ug              |
| <b>Target Name:</b>       | TNFRSF17, CD269, BCM, BCMA |
| <b>Regulatory Status:</b> | RUO                        |

### PRODUCT DETAILS

---

|                               |                                                                                                                                                                |
|-------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Application:</b>           | Flow Cytometry                                                                                                                                                 |
| <b>Format:</b>                | Liquid, PE                                                                                                                                                     |
| <b>Expression Host:</b>       | HEK293                                                                                                                                                         |
| <b>Species:</b>               | Human                                                                                                                                                          |
| <b>Sources:</b>               | Human BCMA protein (NP_001183.2) (Met1-Ala54) with C-terminus Human IgG1 Fc tag is expressed in HEK293 cells and conjugated to PE.                             |
| <b>Accession Number:</b>      | Q02223                                                                                                                                                         |
| <b>Molecular Weight:</b>      | The protein has a predicted molecular weight of 33kDa. Under DTT-reducing conditions, it migrates at approximately 35-45 kDa on SDS-PAGE prior to conjugation. |
| <b>Affinity Tag:</b>          | C-Fc                                                                                                                                                           |
| <b>Formulation:</b>           | 1xPBS buffer, pH7.4, 0.09% NaN3 with a carrier protein                                                                                                         |
| <b>Endotoxin level:</b>       | Not tested                                                                                                                                                     |
| <b>Protein Concentration:</b> | 25µg size is bottled at 0.1mg/mL concentration. 100 µg size is bottled at lot specific concentration.                                                          |
| <b>Storage and Handling:</b>  | Briefly centrifuge the vial upon receipt. An unopened vial may be stored at 2-8°C for up to six months.                                                        |

### BACKGROUND INFORMATION

---

B-cell maturation antigen (BCMA), also known as CD269 or TNFRSF17, is a transmembrane glycoprotein that serves as a critical regulator of B cell development and function. It belongs to the tumor necrosis factor receptor (TNFR) superfamily and is predominantly expressed on plasma cells and a subset of late-stage B cells. BCMA's primary biological role is to promote the survival, differentiation, and long-term maintenance of antibody-producing plasma cells by mediating signals from specific ligands in the TNF family.

Structurally, BCMA is a type I transmembrane protein consisting of an extracellular cysteine-rich domain responsible for ligand binding, a single transmembrane region, and a short cytoplasmic tail that interacts with intracellular signaling molecules. The cytoplasmic domain lacks death domains but recruits TRAF (TNF receptor-associated factor) adaptor proteins to activate downstream pathways such as NF-κB and MAPK. These signaling cascades enhance plasma cell survival and immunoglobulin production, contributing to humoral immunity maintenance. Soluble BCMA, generated through proteolytic cleavage by γ-secretase,


can act as a decoy receptor to regulate ligand availability in the serum.

BCMA binds two main ligands: B-cell activating factor (BAFF, also known as BLyS) and a proliferation-inducing ligand (APRIL). Both ligands are produced by myeloid and stromal cells and support B cell homeostasis. Among these, APRIL binds BCMA with higher affinity and is the primary mediator of BCMA-dependent signaling in plasma cells. The BAFF/APRIL-BCMA axis thus serves as a crucial checkpoint for sustained antibody production and plasma cell survival.


In disease contexts, BCMA is strongly implicated in multiple myeloma (MM) and certain B-cell lymphomas. Its selective overexpression on malignant plasma cells makes it an important diagnostic marker and a therapeutic target. BCMA-directed treatments have revolutionized therapy for multiple myeloma, including antibody-drug conjugates (e.g., belantamab mafodotin), bispecific T cell engagers (e.g., teclistamab, elranatamab), and CAR-T cell therapies (e.g., idecabtagene vicleucel, ciltacabtagene autoleucel). These agents exploit BCMA's restricted expression pattern to deliver targeted cytotoxicity, leading to durable responses in refractory disease. Consequently, BCMA has emerged as a prototypical target in the development of next-generation immunotherapies for hematologic malignancies.

## PRODUCT DATA

**A: BCMA CAR-transfected**  
**Stained with PE-BCMA-Fc**



**B: Mock-transfected**  
**Stained with PE-BCMA-Fc**



CHO cells transfected with either BCMA CAR or Mock plasmid were stained with PE conjugated BCMA (C-Fc) protein at 4ug\_test

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.