

Recombinant Human M-CSF Protein

Catalog Number:	630501, 630502
Size:	20 µg, 100 µg
Target Name:	MCSF, CSF1, CSF-1
Regulatory Status:	RUO

PRODUCT DETAILS

Application:	Bioassay
Format:	Lyophilized from sterile PBS, pH 7.4.
Expression Host:	HEK293
Species:	Human
accession number:	P09603-1
Sources:	A DNA sequence encoding the human CSF1 (P09603-1) (Met1-Asn190) was expressed with an initial Met.
Molecular Weight:	The recombinant human CSF1 consists 158 amino acids and predicts a molecular mass of 18.4 kDa.
Affinity Tag:	None
Purity:	> 85 % as determined by SDS-PAGE.
Endotoxin level:	
Protein Concentration:	Lyophilized
Storage and Handling:	Proteins are stable for up to twelve months from date of receipt at -20°C to -80°C. Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

BACKGROUND INFORMATION

Human macrophage colony-stimulating factor (M-CSF), also known as CSF-1, is a key cytokine that regulates the survival, proliferation, and differentiation of monocytes, macrophages, and osteoclasts. It is produced by a variety of cell types, including fibroblasts, endothelial cells, and stromal cells, and plays a central role in innate immunity, tissue homeostasis, and bone remodeling. M-CSF is essential for the development of the mononuclear phagocyte system and influences macrophage polarization and function in response to environmental cues.

Structurally, M-CSF exists as a homodimeric glycoprotein and can be found in secreted, membrane-bound, and proteoglycan-associated forms generated through alternative splicing. Its primary receptor is CSF-1R (CD115), a class III receptor tyrosine kinase expressed on monocytes and macrophages. Binding of M-CSF induces receptor dimerization and autophosphorylation, activating downstream signaling pathways such as PI3K/AKT, MAPK, and ERK, which promote cell survival and differentiation. M-CSF shares its receptor with another ligand, IL-34, which can also activate CSF-1R and modulate similar biological

processes.

Dysregulation of M-CSF signaling is implicated in various diseases, including cancer, where it promotes tumor-associated macrophage development and immune suppression, as well as in inflammatory conditions and bone disorders such as osteoporosis. Therapeutically, targeting the M-CSF/CSF-1R axis with inhibitors or antibodies is being explored to modulate macrophage activity in cancer and chronic inflammation. Conversely, recombinant M-CSF has potential applications in enhancing immune recovery and tissue repair.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.