

## iF647 Human Annexin V Protein

<b>Catalog Number:</b>	604211, 604212
<b>Size:</b>	25 tests, 100 tests
<b>Target Name:</b>	Annexin A5
<b>Regulatory Status:</b>	RUO

### PRODUCT DETAILS

---

<b>Application:</b>	Flow Cytometry
<b>Format:</b>	iFlour 647
<b>Expression Host:</b>	E.coli
<b>Species:</b>	Human
<b>accession number:</b>	P08758
<b>Sources:</b>	Recombinant Human Annexin A5 (Met1-Asp320) with N-His-Xa tag is expressed in E.coli system. His tag is cut by Xa after purification. This protein is then conjugated to iF647.
<b>Molecular Weight:</b>	This protein has a predicted molecular weight of 35.9 kDa. Under DTT-reducing conditions, the protein migrates at approximately 35 kDa on SDS-PAGE prior to conjugation.
<b>Affinity Tag:</b>	None
<b>Formulation:</b>	1xPBS buffer, pH7.4, 0.09% NaN3 with a carrier protein
<b>Protein Concentration:</b>	Supplied at a lot-specific concentration.
<b>Storage and Handling:</b>	Briefly centrifuge the vial upon receipt. An unopened vial may be stored at 2-8°C for up to six months.
<b>Recommended Usage:</b>	For flow cytometric staining, it is recommended to use 5 uL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. iF647 has an excitation max at 656 nm and an emission max at 670 nm.
<b>Excitation Laser:</b>	Red Laser (633 nm)

### BACKGROUND INFORMATION

---

Annexin A5 is a member of the annexin family, a group of proteins that bind negatively charged phospholipids in a calcium-dependent manner. During apoptosis, phosphatidylserine (PS) translocates to the outer leaflet of the plasma membrane. Annexin A5 specifically binds to this exposed PS, making it a widely used marker for early apoptosis. It is commonly used in combination with propidium iodide (PI) in flow cytometry or microscopy assays